

Thalassemia in the 21st century: Challenges, and solutions, a review article

Suad M Jatal^{1*}; Sawsan J Harfouch¹; Rim M Harfouch²

¹*Faculty of Pharmacy, Alsham Private University ASPU, Latakia, Syria.*

²*Faculty of Pharmacy, Latakia (Tishreen) University, Latakia, Syria.*

***Corresponding author: Suad M Jatal**

Faculty of Pharmacy, Alsham Private University ASPU, Latakia, Syria.

Email: s.j.foph.lat@aspu.edu.sy

Received: Aug 14, 2025; **Accepted:** Sep 12, 2025;

Published: Sep 19, 2025

Citation: Jatal SM, Harfouch SJ, Harfouch RM. Thalassemia in the 21st century: Challenges, and solutions, a review article. *Ann Case Rep Med Images.* 2025; 2(2): 1040.

Introduction

Thalassemia represents a group of inherited disorders characterized by defective hemoglobin synthesis, leading to ineffective erythropoiesis and hemolytic anemia. The disease is most prevalent in the Mediterranean, Middle East, Indian sub-continent, and parts of Southeast Asia and Africa. It is caused by mutations that affect the production of either alpha (α) or beta (β) globin chains, resulting in alpha- or beta-thalassemia, respectively [1].

Genetic basis and pathophysiology

Thalassemia is typically inherited in an autosomal recessive manner. Beta-thalassemia results from mutations in the HBB gene on chromosome 11, leading to reduced (β^+) or absent (β^0) production of beta-globin chains. In alpha-thalassemia, gene deletions in HBA1 and HBA2 on chromosome 16 impair alpha-chain production.

The imbalance between alpha and beta chains causes ineffective erythropoiesis and hemolysis. In beta-thalassemia major, the complete absence of beta-globin leads to severe anemia that manifests in infancy and necessitates lifelong transfusion therapy [2].

Clinical features

Thalassemia presents with a broad clinical spectrum:

- Thalassemia major: Severe transfusion-dependent anemia, hepatosplenomegaly, bone deformities, growth retardation, and iron overload complications.
- Thalassemia intermedia: Moderate anemia, variable transfusion needs, and fewer complications.
- Thalassemia minor: Usually asymptomatic with mild microcytic anemia.
- Complications include cardiac failure, endocrine dysfunction, osteoporosis, and increased susceptibility to infections, particularly in patients who have undergone splenectomy [3].

Diagnosis

The diagnostic approach includes:

1. Complete Blood Count (CBC): Shows microcytic, hypochromic anemia with low MCV and MCH [4].
2. Peripheral Blood Smear: Reveals target cells, anisopoikilocytosis, and nucleated red cells.

3. Hemoglobin Electrophoresis or HPLC:

- o Beta-thalassemia major: High HbF, absent or low HbA.
- o Beta-thalassemia trait: Elevated HbA2 (>3.5%).

4. Genetic Testing: Confirms mutation type; useful for prenatal diagnosis and family screening.

5. *Serum Ferritin and MRI: Evaluate iron overload [5,6].

Management

1. Blood transfusions

- Essential for patients with thalassemia major which should be performed in specialized centers.
- Aims to maintain Hb levels >9–10 g/dL.
- Risks: iron overload, alloimmunization, transmission of infections.

2. Iron chelation therapy

Iron overload from transfusions is managed using:

- Deferoxamine: Subcutaneous infusion.
- Deferiprone and Deferasirox: Oral agents with better compliance.

Chelation is guided by serum ferritin and organ-specific iron measurements [7].

3. Splenectomy

Considered for patients with hypersplenism or high transfusion requirements. It increases infection risk, necessitating vaccination against pneumococcus, meningococcus, and *Haemophilus influenzae* type B [8,9].

4. Hematopoietic Stem Cell Transplantation (HSCT)

- The only curative option.
- Best results in young patients with HLA-matched sibling donors [10].

5. Emerging therapies

- Gene therapy: Lentiviral vector-mediated transfer of functional beta-globin genes has shown success in trials [11].
- Luspatercept: Promotes late-stage erythroid maturation; reduces transfusion burden in transfusion-dependent patients [12].

Prevention and screening

In high-prevalence areas, prevention strategies are critical:

- Carrier screening and genetic counseling.
- Premarital screening programs (e.g., in Cyprus and Iran) have significantly reduced disease incidence.
- Prenatal diagnosis via chorionic villus sampling or amniocentesis [13].

Community education plays a vital role in promoting these programs, especially in regions where consanguinity is common.

Complications

Without adequate management, thalassemia may result in:

- Iron overload: Cardiac dysfunction, hepatic fibrosis, diabetes, hypothyroidism.
- Infections: Particularly post-splenectomy.
- Skeletal deformities: Due to bone marrow expansion.
- Endocrine disorders: Including hypogonadism, delayed puberty, and growth failure [14].

Conclusion

Thalassemia remains a significant global health issue, especially in low-resource settings. While advances in chelation therapy, supportive care, and curative treatments like HSCT and gene therapy have improved survival and quality of life, prevention through carrier screening and public education remains the most cost-effective strategy. Coordinated care involving hematologists, endocrinologists, and infectious disease specialists is vital for optimal outcomes.

References

1. Weatherall DJ, Clegg JB. The Thalassemia Syndromes. 4th ed. Blackwell Science; 2001.
2. Higgs DR, Engel JD, Stamatoyannopoulos G. Thalassemia. *Lancet*. 2012; 379: 373–383.
3. Rund D, Rachmilewitz E. Beta-thalassemia. *N Engl J Med*. 2005; 353: 1135–1146.
4. Sukkariyeh S, Latouf C, Harfouch R. Iron Deficiency Among a Group Syrian Woman: Causes and Treatment. *J Skeleton System*. 2023; 2: 2836–2284.
5. Cappellini MD, Cohen A, Porter J, et al. Guidelines for the Management of Transfusion Dependent Thalassemia (TDT). 3rd ed. Thalassemia International Federation; 2014.
6. Wood JC. Use of magnetic resonance imaging to monitor iron overload. *Hematol Oncol Clin North Am*. 2014; 28: 747–764.
7. Kwiatkowski JL. Management of transfusional iron overload. *J Blood Med*. 2011; 2: 135–149.
8. Davies JM, Barnes R, Milligan D. Update of guidelines for the prevention and treatment of infection in patients with an absent or dysfunctional spleen. *Clin Med (Lond)*. 2002; 2: 440–443.
9. Jatal MJ, Harfouch JS, Harfouch RM. Insights into the Clinical Features of Thalassemia Patients from Latakia, Syria. 2025.
10. Angelucci E, Matthes-Martin S, Baronciani D, et al. Hematopoietic stem cell transplantation in thalassemia major and sickle cell disease: indications and management recommendations from an international expert panel. *Haematologica*. 2014; 99: 811–820.
11. Thompson AA, Walters MC, Kwiatkowski J, et al. Gene Therapy in Patients with Transfusion-Dependent β -Thalassemia. *N Engl J Med*. 2018; 378: 1479–1493.
12. Cappellini MD, Taher AT, et al. A Phase 3 Trial of Luspatercept in Patients with Transfusion-Dependent β -Thalassemia. *N Engl J Med*. 2020; 382: 1219–1231.
13. Al Arrayed S. Campaign to control genetic blood diseases in Bahrain. *Community Genet*. 2005; 8: 52–55.

14. Borgna-Pignatti C, Galanello R. Thalassemias and related disorders: quantitative disorders of hemoglobin synthesis. In: Orkin SH, Nathan DG, Ginsburg D, Look AT, Fisher DE, Lux SE, editors. Nathan and Oski's Hematology of Infancy and Childhood. 7th ed. Saunders. 2009: 1011–1048.